


Gerire Hills

ETHIPA0015



Country: **Ethiopia**

Administrative region: **Somali (Regional State)**

Central co-ordinates: 5.85316 N, 42.10909 E

Area: 3924.57km²

Qualifying IPA criteria

A(i), A(iii)

IPA assessment rationale

The Gerire Hills qualify as an IPA under criterion A(i) as it contains the only known populations of three globally threatened species: *Croton elkerensis* (VU), *Equilabium spananthum* (CR) and *Gnidia elkerensis* (VU), and one of two known populations of *Euphorbia omariana* (VU). Additionally, this site is triggered under criterion A(iii) as it contains the only known distribution of four highly range restricted endemic plants which may be threatened including: *Aloe jacksonii* (DD), *Anacampseros specksii* (DD), *Blepharispermum obovatum* (not assessed) and *Euphorbia bertemariae* (DD).

Site description

The Gerire Hills, which reach a height of ca. 1300 m at Mt Elliot, are located between the Weyb and Wabi Shebelle Rivers in the western Ogaden region of southeastern Ethiopia. Administratively, the majority of this IPA belongs to the Somali Regional State, however a

small portion of the northernmost boundary of the IPA is contained within the Oromia Regional State. The Gerire Hills are part of a southeastern extension of the Ethiopian highlands called the Audo Mountain Range, which consists of the Bur Uare, Monte Elliot, Bur Caddas and Bur Dami plateaus.

This IPA occurs within the Bale floristic region and is a site of botanical significance due to the presence of several range restricted endemic species.

The IPA boundary is defined by topography and geology that has been inferred from satellite imagery and DEMs (GoogleEarth 2023; Landsat GeoCover 1990) and has been delineated to include plateaus and escarpments (850-1300 m) but not the surrounding plains (

Botanical significance

This IPA is a botanically significant site as it contains the entire known populations of eight endemic plant species, namely: *Aloe elkeriana*, *Aloe jacksonii*, *Anacampseros specksii*, *Blepharispermum obovatum*, *Croton elkerensis*, *Equilabium spananthum*, *Euphorbia bertemariae* and *Gnidia elkerensis*. *Aloe elkeriana* and *Aloe jacksonii* have been described from only this locality, however in recent targeted surveys *A. jacksonii* was not found (Friis et al., 2019). The Ethiopian endemic succulent, *E. omariana*, also occurs in this IPA as well as one other location, namely the Sof Omar Caves.

Of these endemics, the following have been assessed: *Aloe elkeriana* (LC), *Aloe jacksonii* (DD), *Anacampseros specksii* (DD), *Croton elkerensis* (VU), *Equilabium spananthum* (CR), *Euphorbia bertemariaea* (DD), *Euphorbia omariana* (EN) and *Gnidia elkerensis* (VU). *Blepharispermum obovatum* has not yet been assessed.

The assessment for the endemic species, *Aloe elkeriana* (LC) is out of date. As it is currently known from only one location and has an estimated area of occurrence of 4km², this species may now be threatened.

This region of Ethiopia is under-collected (Friis et al., 2018), and the gypsum outcrops of southeast Ethiopia in particular have high potential for new species discovery (M. Gilbert, pers. comm., 2023). Seven species at the site concentrated on Burr Caddas plateau are narrow-range endemics currently only known to occur within the IPA boundary, however similar habitat is found over the entire Audo Mountains. Further fieldwork is needed to ascertain whether additional subpopulations of these species occur elsewhere within and outside the Gerire Hills IPA.

Transitional semi evergreen bushland (TSE) is the vegetation community that dominates on the Monte Elliot and the Bur Caddas plateaus. The TSE present within the IPA is a significant outlier from the rest of its mapped distribution and it is highly threatened vegetation type within Ethiopia (van Breugel et al., 2016).

Habitat and geology

The Gerire Hills are characterized by two broad vegetation types: Transitional semi-evergreen bushland (TSE) which is found at ca. 1150 m at higher elevations on the plateau, and *Acacia-Commiphora* bushland which is found on the lower slopes and adjacent plains (Friis et al., 2018). The presence of TSE is clearly indicated by evergreen species such as by a scattered canopy of *Terminalia polycarpa* and *T. haract*, while the *Acacia-Commiphora* plant community is characterized by deciduous trees and shrubs as well as many species displaying succulent traits (Friis et al. 2018). A detailed vegetation description for the Gerire Hills is provided by Friis et al. (2018). The TSE of the Audo Ranges represents an outlier of this vegetation community as it is ca. 160 km from continuous blocks of this vegetation type, which is primarily distributed along the eastern escarpment of the central Ethiopian highlands (Friis et al., 2018). Some of the vegetation present at the Gerire Hills is secondary, and there is strong evidence that the botanical composition of the vegetation has changed significantly over the last ca. 100 years. Species that were recorded by Mogadisco as being dominant at the site such as *Juniperus procera* (LC) and *Buxus hildebrandtii* (not assessed) are no longer so (Friis et al., 2010; Friis et al., 2018).

The geology of the Gerire Hills is unique and highly stratified. The highest points on the Bur Caddas and Monte Elliot plateaus consist of Paleocene sandstones of the Yesomma Formation (Mège et al.

2015). Below the thick sandstone layers are cretaceous limestone layers of the Gorrahei Formation (Mège et al. 2015; Friis et al. 2018; Friis et al 2019). On the northernmost aspects of the mountain range, rocks are significantly tilted exposing underlying gypsum and there are occasional outcrops of gypsum along all edges of the sandstone plateau (D. Mège, pers. comm., 2023). Mège et al. (2015) have demonstrated that erosion by underground rivers and streams and gravity have played a very important role in the creation of the present geomorphology. The tilted rocks and rugged terrain present at the site is caused by the higher rates of erosion of the Gorrahei evaporites which underlay the Yesomma sandstone (Mège et al. 2015). The soil on the sandstone plateau is red and very fine-grained sand, and soils derived from the Gorrahei formation with a high gypsum content are predominately yermosols which are not suitable for agriculture (D. Mège, pers. comm., 2023; Mège et al., 2015). A geomorphological map of the Audo Range can be found in Mège et al. (2015).

This IPA falls within the kolla Ethiopian climactic zone which is a warm to hot semiarid climate (Mège et al., 2015; Asefa et al., 2020). This zone is between 500-1500 m a.s.l. and has an average annual rainfall between 500-800 mm (Mège et al., 2015). Locally the Audo Mountains experience cooler, wetter conditions than the surrounding plains.

Conservation issues

Despite the high endemism of plant species in southeastern Ethiopia, this region is under-represented in the protected area network. Currently the Gerire IPA is not formally protected.

Insufficient data on the flora of southeastern Ethiopia in general is inhibiting effective conservation of Ethiopia's unique and rich plant diversity. The Ogaden has not been better studied to date as a product of a combination of factors; including the remoteness, difficult terrain, limited infrastructure and difficult security conditions, and these issues continue to persist (Mège et al., 2015).

Due to the high amount of gypsum present at the site, this area may be a potential site of future development for gypsum mining. Gypsum is primarily extracted by surface mining, which is inevitably a destructive process for the vegetation cover on the gypsum deposit. While there are not currently known plans to mine the Audo Mountains, mining concessions have been awarded to a company named 'Best Gypsum Industries' to explore deposits in the Somali Regional State (Berhane, 2019) and there is currently some gypsum mining activity within the Oromia Regional State (Plaza-Toledo, 2018).

The Audo Mountains are desirable farmland in an otherwise dry area. For this reason, local communities have favoured the Gerire Hills, where they have practiced agriculture on the plateau, the wider valleys and on the less-steep slopes of this area for centuries (Mogadisco, 1938; translated by Friis et al., 2019). While some

species found on the plateau are likely tolerant of some level of disturbance, ongoing and increased grazing will likely impact survival and recruitment of native plant species. In particular, the TSE which occurs on the plateau is under threat of fragmentation, habitat degradation and deforestation for farming and grazing. This pressure is compounded with a shift from traditional farming practices to modern, more intensive agricultural methods (Friis et al., 2018). Intensification of activity in the area is likely, with the development of a new road connecting El Kere to Imi in the north and Melka Chireti in the south facilitating both the expansion of El Kere and the transportation of large agricultural machinery. This could result in a drastic reduction in habitat for the endemic species for which this IPA has been designated (Friis et al., 2018; Friis, 2019).

Competition from invasive species also poses a threat to the persistence of native plants in the area. The invasive, though native, species *Dodonaea viscosa* subsp. *angustifolia* has been observed to occur within the IPA (Friis et al., 2018). *D. viscosa* subsp. *angustifolia* is an indicator of grazing pressure and former cultivation and outcompetes other native species in these altered environments (Friis et al., 2018).

Inappropriate fire regimes are a threat to the Gerire Hills IPA. Fire is a significant factor shaping the distribution of vegetation in Ethiopia and it acts like a filter for species persistence (van Breugel et al., 2016). Human influence has strongly influenced fire occurrence in Ethiopia (Harris et al., 2023). Human disturbance has altered the fire regime at the site, which has caused a shift in the floristic composition (Friis et al., 2018). In the second half of the 19th century a human-induced fire escaped control on the Monte Elliot and Bur Cadas plateaus and destroyed much of the vegetation there and in the high valleys (Mogadiscio, 1938 translated in Friis et al., 2019). *Juniperus procera* was previously recorded as a dominant at the site; however, recent survey found only a few individuals of *J. procera* remaining along the forest border and on rocky areas in higher elevations of the forest at the Gerire Hills (Friis et al., 2018). While *J. procera* is known to be susceptible to fire, it is likely that this observed decline is also a result of selective logging practices (Friis et al., 2018). TSE and *Acacia-Commiphora* are vegetation communities which would rarely burn naturally; however, with human disturbance fragmenting and igniting fires, the frequency of fire has increased, and the floristic composition has shifted as a result (van Breugel et al., 2016).

Site assessor(s)

Iain Darbyshire, Royal Botanic Gardens, Kew

Sileshi Nemomissa, Addis Ababa University

Sebsebe Demissew, Addis Ababa University

Ermias Lulekal, Addis Ababa University

Gabriella Hoban, Royal Botanic Gardens, Kew

IPA criterion A species

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Croton elkerensis</i> Friis & M.G.Gilbert	A(i)	✓	✓	✓	✓	—	
<i>Gnidia elkerensis</i> Friis & Sebsebe	A(i)	✓	✓	✓	✓	—	
<i>Euphorbia bertemariae</i> Bisseret & Dioli	A(iii)	✓	✓	✓	✓	—	
<i>Euphorbia omariana</i> M.G.Gilbert	A(i)	—	—	✓	—	—	
<i>Equilabium spananthum</i> (A.J.Paton, Friis & Sebsebe) A.J.Paton	A(i)	✓	✓	✓	✓	—	
<i>Aloe jacksonii</i> Reynolds	A(iii)	✓	✓	✓	✓	—	
<i>Anacampseros specksii</i> Dreher	A(iii)	✓	✓	✓	✓	—	
<i>Blepharispermum obovatum</i> Chiov.	A(iii)	✓	✓	✓	✓	—	

IPA criterion C qualifying habitats

HABITAT	QUALIFYING SUB-CRITERION	≥ 5% OF NATIONAL RESOURCE	≥ 10% OF NATIONAL RESOURCE	1 OF 5 BEST SITES NATIONALLY	AREAL COVERAGE AT SITE
---------	--------------------------	---------------------------	----------------------------	------------------------------	------------------------

General site habitats

GENERAL SITE HABITAT	PERCENT COVERAGE	IMPORTANCE
Shrubland - Subtropical/Tropical Dry Shrubland	—	Unknown
Unknown	—	Unknown

Land use types

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
Agriculture (arable)	—	Major
Agriculture (pastoral)	—	Major

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
Harvesting of wild resources	–	Unknown

Threats

THREAT	SEVERITY	TIMING
Agriculture & aquaculture - Annual & perennial non-timber crops - Small-holder farming	High	Ongoing - increasing
Natural system modifications - Fire & fire suppression	High	Ongoing - trend unknown
Residential & commercial development	Medium	Ongoing - trend unknown
Agriculture & aquaculture - Livestock farming & ranching - Small-holder grazing, ranching or farming	Medium	Ongoing - trend unknown
Invasive & other problematic species, genes & diseases	Medium	Ongoing - trend unknown

Management type

MANAGEMENT TYPE	DESCRIPTION	YEAR STARTED	YEAR FINISHED
No management plan in place		–	–

Bibliography

Friis, I., Sebsebe Demissew, & van Breugel, P. 2010. **Atlas of the Potential Vegetation of Ethiopia.**

Asefa, M., Cao, M., He, Y., Mekonnen, E., Song, X., & Yang, J. 2020. **Ethiopian vegetation types, climate and topography.** *Plant Diversity*, Vol 42, page(s) 302-311

Mège, D., Purcell, P., Pochat, S., & Guidat, T. 2015. 'The Landscape and Landforms of the Ogaden, Southeast Ethiopia.' In Billi, P. **Landscapes and Landforms of Ethiopia.**

Google Earth 2023. **Google Earth Pro 2023.**

Wakgari Kebeta Djigsa 2021. **Towards Socially Responsible Mining Investment in Ethiopia: Imagining a New Moral Economy.**

P. van Breugel, I. Friis, Sebsebe Demissew, Jens-Peter Barnekow Lillesø, Roeland Kindt 2016. **Current and Future Fire Regimes and Their Influence on Natural Vegetation in Ethiopia.** *Ecosystems*, Vol 19, page(s) 369–386

Demissew, S., Friis, I. & Weber, O. 2021. **Diversity and endemism of the flora of Ethiopia and Eritrea: state of knowledge and future perspectives.** *Rend. Fis. Acc. Lincei*, Vol 32, page(s) 675–697

Paulo van Breugel, Ib Friis, Sebsebe Demissew 2016. **The transitional semi-evergreen bushland in Ethiopia: characterization and mapping of its distribution using predictive modelling.** *Applied Vegetation Science*, Vol 19, page(s) 335-367

Friis, I 2019. **Cenni monografici sul paese dei Gherire (Mogadiscio, 1938) – a pioneer work on the Gerire Hills in western Ogaden, south-eastern Ethiopia : – introduced, translated and provided with annotations and a revision of the botanical collections.** *Webbia*, Vol 74 (supplement), page(s) 1-108

Friis, I; Gilbert, M. G.; Weber, O.; Van Breugel, P; Sebsebe, D 2019. **The Gerire Hills, SE Ethiopia: ecology and phytogeographical position of an additional local endemic, Anacampseros specksii (Anacampserotaceae).** *Webbia*, Vol 74, page(s) 185-192

Friis, I 1992. **Forests, and forest trees of northeast tropical Africa. Their natural habitats and distribution patterns in Ethiopia, Djibouti and Somalia.** *Journal of Tropical Ecology*, Vol 9 (pub. Royal Botanic Gardens, Kew), page(s) 43-44

Plaza-Toledo, M 2018. **The Mineral Industry of Ethiopia in 2017-2018.**

Berhane Hailemariam 2023. **The concessionaires will invest 4.3**

billion Br to explore and extract minerals.

Harris, L. B., Taylor, A. H., Habtemariam Kassa, Samson Leta & Powell, B 2023. **Humans and climate modulate fire activity across Ethiopia**. *Fire Ecology*, Vol 19

NASA 2000. **Landsat GeoCover 2000**.

Friis, I., Gilbert, M. G., Paton, A. J., Weber, O., van Breugel, P., & Sebsebe Demissew 2018. **The Gerire Hills, a SE Ethiopian outpost of the transitional semi-evergreen bushland: vegetation, endemism and three new species, *Croton elkerensis* (Euphorbiaceae), *Gnidia elkerensis* (Thymelaeaceae), and *Plectranthus spananthus* (Lamiaceae)**. *Webbia*, Vol 73:2, page(s) 203-223