

# Kasyoha-Kitomi Central Forest Reserve

## UGATIPA7

Country: **Uganda**

Administrative region: **Western (Region)**

Central co-ordinates: **-0.27544 N, 30.23853 E**

Area: **384km<sup>2</sup>**

### Qualifying IPA criteria

A(i), B(ii)

### IPA assessment rationale

Kasyoha-Kitomi is recognised as an IPA under sub-criterion A(i). One Critically Endangered, five Endangered and six Vulnerable species are known from this site. Urgent research is needed to establish if four of these species are still extant at this site, particularly *Diospyros katendei* which is only known from this site globally. With six species range-restricted or endemic species this site also meets sub-criterion B(ii), representing an exceptional number of species of high conservation importance. Finally sub-criterion C(iii) is triggered by the presence of Lake Victoria drier peripheral semi-evergreen Guineo-Congolian rainforest (CR). Kibale encompasses close to 7% of the national resource of this habitat and is one of the five best sites nationally.

### Site description

Kasyoha-Kitomi Central Forest Reserve covers 384 km<sup>2</sup> in area and falls within the districts of Bushenyi, Ibanda and Kamwenge of Uganda's Western Region. Located within the Albertine Rift Area, the site is around 25 km east of Lake Edward and 10 km south of Lake George. The Kyambura River runs through this site, meeting the Kazinga Channel downstream between these two lakes.

The site was surveyed in 2023 as part of fieldwork for the Uganda TIPAs project.

### Botanical significance

Kasyoha-Kitomi is an important site for national and Albertine Rift endemics, many of which are threatened. Of great importance is *Diospyros katendei*, a Critically Endangered tree that is endemic to Kasyoha-Kitomi and is only known from a single individual. Searches have been undertaken to locate this species in recent years but there have been no further individuals found and, as a *Diospyros* species, it is highly likely to have been targeted for its high-quality timber (IUCN SSC East African Plants Red List Authority 2013). Another threatened tree species *Ficus katendei* (EN), similarly named after

the Ugandan botanist Anthony Katende, may have also been extirpated from this site. While this species is found at one other site, Bwindi-Impenetrable National Park, the only known individual at Kasyoha-Kitomi was felled to build a bridge (Gereau et al., In prep.). However, there are large areas of Kasyoha-Kitomi that have yet to be searched for either of these species and, therefore, it is possible that they are still extant at this site. Further research is urgently needed, particularly given the threats to this site, to understand whether there is an opportunity to conserve *D. katendei* and *F. katendei* here; particularly for the former of the two which may be saved from extinction through conservation at Kasyoha-Kitomi.

This IPA is also of great conservation importance for several other threatened species. Kasyoha-Kitomi is, for instance, the only site nationally from which *Aframomum spirologulatum* is known and one of only two protected areas globally in which this species occurs. In addition, another Endangered species *Uvariadendron magnificum*, was described as a "dominant understorey tree" when collected in 1969 (Synott #540) and has been collected more recently in 2017. This species is endemic to Uganda and Kasyoha-Kitomi likely represents the most important site for this species nationally. This IPA is also an important site for *Brachystephanus roseus*, an Albertine Rift endemics known from only two other protected areas in Uganda.

Six Vulnerable species are known from this IPA. One of these species, *Musanga leo-errariae*, was collected during 2023 fieldwork by the Uganda TIPAs team at this site. This tree is an Albertine Rift endemic and occurs occasionally in recently opened gaps in the canopy at Kasyoha-Kitomi. Another Vulnerable species recorded on fieldwork is *Rinorea tshingandaensis*, this species was uncommon at this site and globally only known from Uganda and eastern D.R. Congo (Kalema & Beentje, 2012). Two other Vulnerable species, *Crotalaria adenocarpoides* and *Vernonia parapetersii*, were collected in 1970 on Lubare Ridge in rocky, grassland habitats (Lye #5473, 5480). In recent decades, large areas of this grassland have been replaced by forest plantation and some small-scale agriculture (Google Earth 2023). Some small areas of Lubare Ridge continue to support grassland and so it is of great importance that further surveys are undertaken to ascertain whether these grassland species persist within this IPA.

Vulnerable timber tree *Mimusops bagshawei* was likely observed on fieldwork in 2023 for the Uganda TIPAs project, although it was not possible to collect leaf or fertile material to confirm this determination. This fieldwork did, however, find a new record of *Massularia acuminata* and *Rytigynia bagshawei* var. *lebrunii* for both Uganda and the Flora of Tropical East Africa region. There are likely several more plant species that are of botanical significance at this site. Further research is needed, particularly in harder to access sections, to fully categorise its biodiversity importance.

In addition to species of conservation importance, Kasyoha-Kitomi is also a key site for the conservation of Lake Victoria drier peripheral semi-evergreen Guineo-Congolian rainforest. This habitat has been assessed as Critically Endangered nationally (Richard et al., In review), and is thought to have been more widespread in western Uganda. Kasyoha-Kitomi CFR is one of the best sites nationally for conserving this species.

---

## Habitat and geology

Kasyoha-Kitomi occupies a hilly landscape, reaching a maximum altitude of 2108 m in the northeast, and is geologically complex including quartzites, schists, gneisses, shales and phyllites in the underlying rocks (Howard 1991). The site is dominated by moist forest. Langdale-Brown et al. (1964) classify this forest as mixture of Parinari Forest and Albiza-Markhamia Forest. Our surveys of this site were largely limited to the westerly edges of the reserve within moist, well-developed, secondary forest where species such as *Shirakiopsis* and *Strombosia* were common.

---

## Conservation issues

Kasyoha-Kitomi Central Forest Reserve was established in 1932. With neighbouring protected areas Queen Elizabeth National Park, Kyambura Wildlife Reserve, Kalinzu and Kakasi Central Forest Reserves, Kasyoha-Kitomi forms part of a conservation corridor from Fort Portal to the D.R. Congo border south of Lake Edward (UNEP-WCMC and IUCN 2022).

The site has previously experienced high-levels of disturbance through pitsawing, covering well over half of the reserve area, however, the steep topology has prevented mechanised tree felling (Howard 1991). The site continues to be threatened by gold mining, at the Kitaka deposit in the north-east, with a prospecting license covering 40 km<sup>2</sup> granted to Simba mining group. The mining and associated access roads have clearly led to deforestation at this site (Google Earth 2023; Global Forest Watch 2023). The chairman of Simba group was reported to have claimed that, for every tree cut at the site, 40 more would be planted (Daily Monitor 2013). There are no further details of such a commitment and as such it is unlikely to be binding.

Tea plantations cover large areas of land to the south of the site, landowners and tea estates have previously bought out small-scale farmers leaving them with little land to cultivate (Raben et al. 2007). In turn, shortage of land, combined with high population densities in the area, puts pressure on the habitats of Kasyoha-Kitomi and has led agricultural encroachment and illegal harvesting of timber and charcoal production (Bitariho and Babaasa 2016).

Community Forest Management (CFM) at the site was established by Nature Uganda within the Participatory Environmental Management project between 2007-2011, to curb these threats. Several CFM groups continue to work with site managers. Forest plantations within the site were established outside existing natural forests (BirdLife International 2008). Much of this forestry appears to be on Lubare Ridge, however, satellite imagery shows this area to

have no forest cover in previous decades (Google Earth 2023). In addition, the hilly grasslands of the south of reserve have been given over to small-scale tea plantations, while cattle are grazed in rocky infertile areas (Raben et al. 2007). This loss of habitat has conservation implications for Vulnerable grassland species *Crotalaria adenocarpoides* and *Vernonia parapetersii* (see "Botanical Significance").

---

## Site assessor(s)

Sophie Richards, Royal Botanic Gardens, Kew

Iain Darbyshire, Royal Botanic Gardens, Kew

---

## IPA criterion A species

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Brachystephanus roseus</i> Champl.	A(i)	✓	✓	✓	–	–	Unknown
<i>Dasylepis eggelingii</i> J.B.Gillett	A(i)	✓	–	–	–	–	Occasional
<i>Uvariadendron magnificum</i> Verdc.	A(i)	✓	✓	✓	–	–	Frequent
<i>Vernonia parapetersii</i> C.Jeffrey	A(i)	✓	✓	✓	–	–	Unknown
<i>Diospyros katendei</i> Verdc.	A(i)	✓	✓	✓	✓	–	Scarce
<i>Ficus katendei</i> Verdc.	A(i)	✓	✓	✓	–	–	Unknown
<i>Polystachya hastata</i> Summerh.	A(i)	–	✓	✓	–	–	Unknown
<i>Balsamocitrus dawei</i> Stapf	A(i)	✓	✓	✓	–	–	Unknown
<i>Musanga leoerrerae</i> Hauman & J.Léonard	A(i)	–	✓	✓	–	–	Occasional
<i>Aframomum spiroligulatum</i> Lock & A.D.Poulsen	A(i)	✓	✓	✓	–	–	Unknown
<i>Crotalaria adenocarpoides</i> Taub.	A(i)	✓	–	–	–	–	Unknown
<i>Rinorea tshingandaensis</i> Taton	A(i)	✓	–	✓	–	–	Occasional

## IPA criterion C qualifying habitats

HABITAT	QUALIFYING SUB-CRITERION	≥ 5% OF NATIONAL RESOURCE	≥ 10% OF NATIONAL RESOURCE	1 OF 5 BEST SITES NATIONALLY	AREAL COVERAGE AT SITE
Lake Victoria drier peripheral semi-evergreen Guineo-Congolian rainforest (CR)	C(iii)		–		268.3

## General site habitats

GENERAL SITE HABITAT	PERCENT COVERAGE	IMPORTANCE
Forest - Subtropical/Tropical Moist Lowland Forest	—	Major
Forest - Subtropical/Tropical Moist Montane Forest	—	Major
Grassland - Subtropical/Tropical High Altitude Grassland	—	Minor
Artificial - Terrestrial - Plantations	—	Minor
Artificial - Terrestrial - Arable Land	—	Minor

## Land use types

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
Nature conservation	—	Major
Agriculture (arable)	—	Minor
Forestry	—	Minor

## Threats

THREAT	SEVERITY	TIMING
Energy production & mining - Mining & quarrying	High	Ongoing - increasing
Agriculture & aquaculture - Annual & perennial non-timber crops - Small-holder farming	Medium	Ongoing - trend unknown
Agriculture & aquaculture - Wood & pulp plantations	Medium	Ongoing - trend unknown
Biological resource use - Logging & wood harvesting - Intentional use: subsistence/small scale (species being assessed is the target) [harvest]	Medium	Ongoing - trend unknown

## Protected areas

PROTECTED AREA NAME	PROTECTED AREA TYPE	RELATIONSHIP WITH IPA	AREAL OVERLAP
Kasyoha-Kitomi Central Forest Reserve	Forest Reserve (conservation)	protected/conservation area matches IPA	384

## Conservation designation

DESIGNATION NAME	PROTECTED AREA	RELATIONSHIP WITH IPA	AREAL OVERLAP
Kasyoha-Kitomi Forest Reserve	Important Bird Area	protected/conservation area matches IPA	384
Kasyoha-Kitomi Forest Reserve	Key Biodiversity Area	protected/conservation area matches IPA	384

## Management type

MANAGEMENT TYPE	DESCRIPTION	YEAR STARTED	YEAR FINISHED
Protected Area management plan in place		–	–

## Bibliography

Howard, P. C. 1991. **Nature Conservation in Uganda's Tropical Forest Reserves.**

IUCN & UNEP-WCMC 2022. **The World Database on Protected Areas (WDPA).**

Langdale-Brown, I., Osmaston, H. A., & Wilson, J. G. 1964. **The Vegetation of Uganda and its Bearing on Land-Use.**

IUCN SSC East African Plants Red List Authority 2013. **Diospyros katendei. The IUCN Red List of Threatened Species 2013: e.T35799A47348409..**

Gereau, R.E., Beentje, H.J., Kabuye, C., Luke, W.R.Q., Nshutiyayesu, S. & Ntore, S. In prep.. **Ficus katendei. The IUCN Red List of Threatened Species.**

Google Earth 2023. **Google Earth Pro 2023.**

World Resources Institute 2023. **Global Forest Watch 2023.**

Daily Monitor 2013. **Mining firm breathes new life into Kitaka gold deposits.**

Raben, K., Nyingi, J., Loserian, D., Akello, Z. & Kidoido, M. 2007. **Kasyoha-Kitomi Landscape, Uganda Report: Local Stakeholders' Use of Forest Reserves in Kasyoha-Kitomi Forest Landscape, Uganda And Nguru South Forest Landscape, Tanzania.**

Bitariho, R. & Babaasa, D. 2016. **The Status of Biodiversity in Kasyoha-Kitomi Central Forest Reserve: A Survey Report.**

BirdLife International 2008. **Important Bird Areas factsheet: Kasyoha-Kitomi Forest Reserve.**

Richards, S., Kalema, J., Ojelel, S., Williams, J. & Darbyshire, I. In review. **Improving the application of Important Plant Areas to conserve threatened habitats: a case study of Uganda.**  
Conservation Science and Practice