


# Mbam Minkom Hills

CMNTIPA008



Country: Cameroon

Administrative region: Centre (Region)

Central co-ordinates: 3.96150 N, 11.39150 E

Area: 86km<sup>2</sup>

## Qualifying IPA criteria

A(i)

## IPA assessment rationale

Mbam Minkom Hills qualifies as a potential IPA under criterion A(i) through the presence of around 20–25 threatened species with important populations at the site. Of these, *Coffea fotsoana* and *Aristolochia preussii* have been assessed as Critically Endangered and *Talbotiella bretelerii* (known only from two other small, highly threatened sites in Yaoundé) as Endangered. Mbam Minkom is the only known Cameroon location for *Polystachya carnosa* and *Polystachya kubalae* and may represent the only population of *Coffea fotsoana*, a potentially important coffee wild relative. Given the high number of threatened Orchidaceae, one of the few groups for which the site has been well surveyed, it seems very likely that there are more threatened taxa from other plant families yet to be recorded.

## Site description

The Mbam Minkom Hills are a series of steep prominences located around 15 km northwest of central Yaoundé, the capital city of Cameroon. Rising to 1,295 m, the highest point in Centre region they represent the highest and largest of the several granulitic (metamorphic) inselberg areas in and around the west of the city. Unlike most of the other prominences much of their vegetation is intact, at least in the submontane altitudinal zone (>900 m). The lower slopes at the perimeter of the area are more degraded by cultivation with farmland and farm fallow.

## Botanical significance

Despite their proximity to Yaoundé, the various hills in and around the city have revealed many threatened and rare species of interest. Although relatively little surveyed, Mbam Minkom represents the largest and most intact of these areas and the best hope for conserving some of the species endemic to the Yaoundé area. Simo et al. (2009) have described the orchid flora in recent years but there have been relatively few other surveying efforts. Several of Zenker and Staudt's important historic collections from "Yaoundé Station" do not have a known location and have not been recorded in other parts of the country (Cheek et al., 2010). They probably have their best chance of surviving or being rediscovered in the Mbam Minkom Hills.

## Habitat and geology

Geologically Yaoundé, the Mbam Minkom Hills and most of the southern part of Central province sit within a large meta-sedimentary area of the South Cameroon Plain, south of the Sanaga fault and north of the north-thrusting ancient Congo craton (Ngnotue et al., 2012). These hills are all formed of granulites and migmatites also referred to as embrichite gneiss (Achoundong, 1985), relatively high grade metamorphic rocks formed from sedimentary and igneous protoliths and apparently dating from around 600 mya (Nzenti, 1988; Tchouatcha et al., 2018; Ngnotue et al., 2012). Rainfall measurements are not available for the site itself but precipitation is relatively low in Yaoundé at 1,541 mm per annum (period 1971–2000; World Meteorological Organization, 2021). This is below the level normally thought necessary to sustain evergreen tropical forest (Cheek et al., 2011), although it is likely to vary through the site. The steep western faces intercept westerly monsoon winds, resulting in differences in the physiognomy and composition of the vegetation (Noumi, 2015; Simo et al 2009). The precipitation regime is of the Guinean type with a small (March–June) and greater (September–November) wet season interspersed with a drier period (July–August), and a second more severe dry period between December and February when mean monthly rainfall drops below the relatively flat mean monthly temperature curve (range: 22.8–25.47 °C) on a on a Walter–Leith type chart (Simo et al., 2009; Bissaya et al., 2014; Noumi, 2015). Relative to other montane areas, the Mbam Minkom range may exhibit a negative Massenerhebung (mountain mass elevation) effect due to its relative isolation from other large mountains, potentially resulting in plants having lower altitudinal tolerances and high altitude species occurring at lower altitudes than elsewhere (Noumi, 2015). The vegetation consists of remaining areas of lowland forest much degraded by cultivation, relatively intact submontane forest at 800–1,300 m, and “inselberg” areas characterised by sparse vegetation tolerant of heat and water deficit such as *Microdracoides squamosus* (Droissart et al., 2006; Onana & Cheek, 2011). “Mbam Minkom” means “rocky domes” in the Ewondo language and the steep topology leaves substantial areas of exposed rock. Forest is semi-deciduous on the lower, eastern slopes but evergreen on the summit. There are many rivers and streams beneath the canopy, often forming steep gullies.

## Conservation issues

The proximity of the Mbam Minkom Hills to the capital city presents various conservation challenges and opportunities. Yaoundé was the second largest city in Cameroon at the last census but has been estimated to be growing at over 5% per year and may now have the largest population in the country at over 4 million (United Nations, 2018). It is predicted to reach 5.7 million by 2030. Most of the other hills and areas of vegetation on the western side of the city have been seriously degraded or lost due to small scale agriculture, suburban spread, hotel development, extraction of timber and quarrying. These are all threats to Mbam Minkom as the city enlarges and develops (Nkwemoh et al., 2017, 2018).

Noumi (2015) reports that much forest has been destroyed by small scale agriculture around the edges and on the lower slopes of the Mbam Minkom area but that the interior is largely intact. The western side which is steeper and wetter is less degraded, even though it drops to lower altitudes of around 650 m (Simo et al., 2009). Awa II et al. (2009) report agriculture, logging and settlement within the forest as threats.

The growing size of Yaoundé and its status as a capital city also present an opportunity to benefit the increasing urban population and add prestige to the capital city by protecting the natural heritage of this site. Banco National Park in the city of Abidjan is a possible model. The potential to contribute to clean air, fresh water, leisure, tourism and education present powerful arguments for treating Mbam Minkom and other remaining local inselbergs as integral to the development of the city. Universities and schools utilise Mbam Minkom for fieldwork sites essential for forestry and biodiversity training. The beauty of the landscape is very striking to visitors and there is potential for increased tourism. However, Awa II (2009) report that a tourism drive led to deforestation of important trees and loss of nesting sites to harvest timber for huts at the summit. Unfortunately these huts were poorly sited and were dilapidated by 2009 (O. Lachenaud, 2021, pers. comm. 29 June). Outcomes such as this highlight the importance of integrating biodiversity education and management into well-managed eco-tourism development. The area demarcated here is narrowly separated from other proposed IPAs (Mt Kala, Eloumden and Mt Minloua). There is potential to integrate these sites and create a much more effective conservation area.

---

## Site assessor(s)

Bruce Murphy, Royal Botanic Gardens, Kew

Xander van der Burgt, Royal Botanic Gardens, Kew

Professor Bonaventure Sonke, University of Yaounde I

Dr Vincent Droissart, Institute of Research for Development.  
Marseille

Dr. Roger Fotso, WCS Cameroon

---

## IPA criterion A species

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Angraecum egertonii</i> Rendle (syn. <i>Ancistrorynychus egertonii</i> )	A(i)	✓	✓	✓	–	–	
<i>Bulbophyllum teretifolium</i> Schltr.	A(i)	✓	✓	✓	–	–	
<i>Cheirostylis divina</i> (Guinea) Summerh. var. <i>ochyrae</i> Szlach. & Olszewski	A(i)	✓	✓	✓	–	–	
<i>Coffea fotsoana</i> Stoffelen & Sonké	A(i)	✓	✓	✓	–	–	
<i>Crossandra obanensis</i> Heine	A(i)	✓	✓	✓	–	–	
<i>Diaphanthe bueae</i> (Schltr.) Schltr.	A(i)	✓	✓	–	–	–	
<i>Hymenostegia viridiflora</i> Mackinder & Wieringa	A(i)	✓	✓	✓	–	–	
<i>Polystachya carnosa</i> P.J.Cribb & Podz.	A(iii)	✓	✓	✓	–	–	
<i>Polystachya kubalae</i> Szlach. & Olszewski	A(i)	✓	✓	✓	–	–	
<i>Stolzia grandiflora</i> P.J.Cribb subsp. <i>lejolyana</i> Stévant, Droissart & M.Simo	A(i)	✓	✓	✓	–	–	
<i>Talbotiella breтели</i> (Aubrév.) Mackinder & Wieringa	A(i)	✓	✓	✓	–	–	
<i>Ardisia ototomoensis</i> Taton	A(i)	✓	✓	✓	–	–	
<i>Pterorhachis zenkeri</i> Harms	A(i)	–	–	–	–	–	
<i>Boutiquea platypetala</i> (Engl. & Diels) Le Thomas	A(i)	✓	✓	–	–	–	

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Aristolochia preussii</i> Engl.	A(i)	✓	✓	✓	–	–	
<i>Chassalia laikomensis</i> Cheek	A(i), A(iii)	✓	–	–	–	–	
<i>Mussaenda epiphytica</i> Cheek	A(i)	✓	✓	✓	–	–	
<i>Cola hypochrysea</i> K.Schum.	A(i), A(iii)	✓	✓	✓	–	–	
<i>Khaya ivorensis</i> A.Chev.	A(i)	–	–	–	–	✓	
<i>Garcinia kola</i> Heckel	A(i)	–	–	–	–	✓	
<i>Entandrophragma angolense</i> (Welw.) C.DC.	A(i)	–	–	–	–	✓	
<i>Sabicea laxa</i> Wernham	A(i), A(iii)	✓	✓	✓	–	–	
<i>Polystachya kornasiana</i> Szlach. & Olszewski	A(i)	✓	✓	✓	–	–	
<i>Psychotria villicarpa</i> O.Lachenaud subsp. <i>villicarpa</i>	A(i)	✓	✓	✓	–	–	
<i>Chassalia manningii</i> O.Lachenaud ined.	A(i)	✓	✓	✓	–	–	
<i>Psychotria senterrei</i> O.Lachenaud	A(i)	✓	✓	✓	–	–	
<i>Calycosiphonia macrochlamys</i> (K.Schum.) Robbr.	A(i)	–	–	–	–	–	
<i>Kylicanthe cornuata</i> Descourv. & Stévant & Droissart	A(i)	✓	✓	✓	–	–	
<i>Oxyanthus doucetii</i> Sonké & O.Lachenaud	A(i)	✓	✓	✓	–	–	
<i>Ardisia koupensis</i> Taton	A(i)	✓	✓	✓	–	–	
<i>Hymenocoleus glaber</i> Robbr.	A(i)	✓	–	–	–	–	
<i>Psychotria camerunensis</i> E.M.A.Petit	A(i)	–	–	✓	–	–	

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
---------	--------------------------	---------------------------	-----------------------------	------------------------------	--------------------------	------------------------------	-------------------

## IPA criterion C qualifying habitats

HABITAT	QUALIFYING SUB-CRITERION	≥ 5% OF NATIONAL RESOURCE	≥ 10% OF NATIONAL RESOURCE	1 OF 5 BEST SITES NATIONALLY	AREAL COVERAGE AT SITE
---------	--------------------------	---------------------------	----------------------------	------------------------------	------------------------

## General site habitats

GENERAL SITE HABITAT	PERCENT COVERAGE	IMPORTANCE
----------------------	------------------	------------

## Land use types

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
---------------	------------------	------------

## Threats

THREAT	SEVERITY	TIMING
Geological events - Avalanches/landslides	Medium	Past, likely to return
Residential & commercial development - Housing & urban areas	Medium	Ongoing - increasing
Energy production & mining - Mining & quarrying	Medium	Future - inferred threat
Agriculture & aquaculture - Annual & perennial non-timber crops - Shifting agriculture	High	Ongoing - trend unknown
Biological resource use - Logging & wood harvesting - Unintentional effects: subsistence/small scale (species being assessed is not the target) [harvest]	High	Ongoing - trend unknown
Natural system modifications - Fire & fire suppression - Increase in fire frequency/intensity	High	Ongoing - trend unknown

## Conservation designation

DESIGNATION NAME	PROTECTED AREA	RELATIONSHIP WITH IPA	AREAL OVERLAP
Mbam Minkom-Mt Kala IBA	Important Bird Area	protected/conservation area encompasses IPA	—

## Management type

MANAGEMENT TYPE	DESCRIPTION	YEAR STARTED	YEAR FINISHED
No management plan in place		—	—

## Bibliography

- Onana J.-M. & Cheek M. 2011. **Red Data Book of the flowering plants of Cameroon.**
- Cheek, M., Harvey, Y. & Onana, J.M. 2011. **The Plants of Mefou Proposed National Park, Yaoundé, Cameroon.**
- BirdLife International 2020. **Important Bird Areas factsheet: Mbam Minkom - Kala.**
- Zogning, A., Ngouanet, C. & Tiafack, O. 2007. **The catastrophic geomorphological processes in humid tropical Africa: A case study of the recent landslide disasters in Cameroon.** *Sedimentary Geology*, Vol 199, page(s) 13 – 27
- Bissaya, R., Ghogomu, R.T., Moundi, A., Njom, B. & N.S. Kanouo 2014. **Utilisation des données géologiques et gestion des informations multi-sources pour l'analyse de l'aléa glissement de terrain/éboulement dans le secteur Nord-Ouest de la région de Yaoundé.** *Afrique SCIENCE*, Vol 10(3), page(s) 113 - 133
- Nkwemoh, C.A.; Tchindjang, M. 2018. **Urban sprawl and agriculture: A case study of the Yaounde metropolis (Cameroon).** *Revue Scientifique et Technique Forêt et Environnement du Bassin du Congo*, Vol 10, page(s) 45-58.
- Nkwemoh, C.A., Tchindjang, M. & Afungang, R.N. 2017. **The Impact of Urbanization on the Vegetation of Yaounde, (Cameroon).** *International Journal of Innovative Research & Development*, Vol 6(5), page(s) 6-18
- Achoundong, G. 1996. **Les forêts sommitales du Cameroun: Végétation et flore des Collines de Yaoundé.** *Bois et forêt des tropiques*, Vol 247, page(s) 37-52
- Noumi, E. 2015. **Floristic structure and diversity of a tropical sub-montane evergreen forest, in the Mbam minkom massif (Western Yaoundé).** *Journal of Biology and Life Science*, Vol 6(1), page(s) 149-193
- Simo, M., Droissart, V., Sonké, B. & Stévant, T. 2009. **The Orchid Flora of the Mbam Minkom Hills (Yaoundé, Cameroon).** *Belgian Journal of Botany*, Vol 142(2), page(s) 111-123
- Nzenti, J. P. 1987. **Pétrogenèse des migmatites de Yaoundé (Cameroun). Élément pour un modèle géodynamique de la chaîne pan africaine Nord Equatoriale.** Thèse Doctorat. (pub. Univ. Nancy I), page(s) 147
- Nzenti, J.P., Barbey, P., Macaudiere, J. & Soba, D. 1988. **Origin and evolution of the late Precambrian high-grade Yaounde gneisses (Cameroon).** *Precambrian Research*, Vol 38, page(s) 91-109
- Tchouatcha, M.S., Kouske, A.P., Njiosseu, E.L.T., Ngouem, P.A., Ngotue, T., Njinchuki, D.N. & Nzenti, J.P. 2018. **Preserved Sedimentary Features in the Pan-African High-Grade Metamorphic Rocks from the Yaoundé Series (Cameroon).** *Journal of Geosciences and Geomatics*, Vol 6(3), page(s) 94-102
- Ngotué, T., Ganno, S., Nzenti, J.P., Schulz, B., Tchaptchet T.D. & Suh, C.E. 2012. **Geochemistry and geochronology of Peraluminous High-K Granitic Leucosomes of Yaoundé Series ) Cameroon. Evidence for a Unique Pan-African Magmatism and Melting Event in North Equatorial Fold Belt.** *International Journal of Geosciences*, Vol 3, page(s) 525-548
- Awa II, T., Dzikouk, G. & Norris, K. 2009. **Breeding distribution and population decline of globally threatened Grey-necked Picathartes Picathartes oreas in Mbam Minkom Mountain Forest, southern Cameroon.** *Bird Conservation International*, Vol 19, page(s) 254–264
- Droissart, V., Sonké, B. & Stévant, T. 2006. **Les Orchidaceae endémiques d'Afrique centrale atlantique présentes au Cameroun.** *Systematics and Geography of Plants*, Vol 76(1)
- United Nations, Department of Economic and Social Affairs, Population Division 2018. **World Urbanization Prospects: The 2018 Revision, Online Edition.**