


Mount Karkarha

ETHIPA007



Country: **Ethiopia**
 Administrative region: **Southern Nations, Nationalities, and Peoples' (Regional State)**
 Central co-ordinates: **6.96532 N, 35.67209 E**
 Area: **138km²**

Qualifying IPA criteria

A(i)

IPA assessment rationale

Mount Karkarha qualifies as an IPA under criterion A(i) due to the presence of the globally threatened, Ethiopian endemic species *Dorstenia soerenseii* Friis (EN) and *Scadoxus nutans* (VU) for which this IPA is considered to be an important site.

Site description

Mount Karkarha, also known as Kaka mountain, is located within the Southern Nations, Nationalities, and People's Regional State, in the Kefa floristic region. It is not to be confused with Mount Kaka in the Oromia Regional State. Some sources refer to the IPA as Gem mountain, however this name is unconfirmed and requires further research (Kassa et al., 2017a; Kassa et al., 2019). Elevation of the mountain ranges from 1808 m to 2685 m. The Mount Karkarha IPA is situated approximately 50 km west of the Bonga Forests IPA and approximately 17 km SE of the Shako-Bench Forest IPA. It is bordered to the north by the town of Mizan Teferi. A road that joins the B53, the main road between Mizan Teferi and Bonga, runs through the northernmost part of the IPA.

The mountain is under researched in terms of its biodiversity and there has not been a comprehensive botanical survey to date, but it

is known to support intact forest with rare and threatened species present.

Botanical significance

The Mount Karkarha IPA is characterised by Moist Evergreen Afromontane Forest (MAF) (Friis et al., 2010). There has been only very limited botanical surveying of this site to date. However, the IPA appears to hold an extensive area of partially intact MAF forest and is therefore an important example of this threatened habitat.

This IPA is of botanical significance as one of the few localities of the globally threatened, Ethiopian endemic *Dorstenia soerenseii* Friis (EN), an understory shrub often found along small streams and in deep shade. Records of that species are only known from Bonga, Sheka, Mount Karkarha, and near Maji. It also contains a population of a second threatened Ethiopian endemic, the attractive herb *Scadoxus nutans* (Friis & I.Björnstad) Friis & Nordal.

According to Kassa et al. (2018), three Ethiopian endemics are present within the northern slopes of the forest: *Solanecio gigas* (Vatke) C.Jeffrey, *Millettia ferruginea* Hochst. (LC) and *Vernonia tewoldei* Mesfin (LC). Three further Ethiopian endemics were identified by Kassa et al. (2018): *Cirsium schimperi* (Vatke) C. Jeffrey, *Echinops kebericho* Mesfin (NT) and *Pentanema confertiflorum* (A.Rich.) D.Gut.Larr. et al. However, the natural distribution of these latter species does not include the Kefa floristic region (Mesfin Tadesse, 2004) and so their presence at this site is questionable. The Ethiopian endemic *Bidens pachyloma* (Oliv. & Hiern) Cufod., has been recorded within agroforestry areas at the edge of the IPA (Kassa et al., 2018).

Also of note is the widespread medicinal timber species, *Prunus africana* (Hook.f.) Kalkman (VU), reported at low levels within the IPA (Kassa et al., 2018).

Habitat and geology

Mount Karkarha is characterised by steep terrain, dominated by Moist Evergreen Afromontane Forest (Friis et al., 2010). The northern slopes of the IPA are located within the upper Gacheb catchment which drains into the White Nile through the Baro-Akobo river system (Kassa et al., 2017a).

Botanical records from 1976 described the IPA as montane forest with *Aningeria* spp., *Schefflera* spp., and *Ficus* spp. dominating the upper canopy; *Cyathea manniana* Hook. is abundant in the lower canopy; the ground layer is dominated by terrestrial ferns such as *Didymochlaena*, *Tectaria*, *Marratia*, and *Thelypteris* (Gilbert #4203). *Grewia ferruginea* Hochst. ex A.Rich., *Vernonia amygdalina* Delile, and *Solanecio mannii* (Hook.f.) C.Jeffrey are also common within the lower canopy (Kassa et al., 2017b). Epiphytic ferns, such as *Elaphoglossum deckenii* (Kuhn) C.Chr., have been recorded within the IPA (Roux, 2011).

Kassa et al. (2018) recorded the most common tree species within edges of the northern slopes of the IPA as *Juniperus procera* Hochst. ex Endl., *Cyathea manniana*, *Polyscias fulva* (Hiern) Harms, *Astropanax abyssinicus* (Hochst. ex A.Rich.) Seem. (*Schefflera abyssinica*), and the introduced *Grevillea robusta* A.Cunn. ex R.Br. The dominant shrub species were *Vernonia amygdalina*, *Solanecio gigas*, and the introduced *Delonix regia* (Boijer) Raf. The ground flora is dominated by *Plectranthus barbatus* Andrews, *Bidens pilosa* L., and *Veronica persica* Poir..

Within the Gacheb catchment, agroforestry of *Coffea arabica* L. as a cash crop, and *Ensete ventricosum* Welw. Cheesman and *Aframomum corrorima* Braun as food crops is common (Kassa et al., 2019). Maize is the most common crop grown in the agricultural fields surrounding the IPA (Kassa et al., 2019). Native trees such as *Albizia gummifera* J.F.Gmel C.A.Sm., *Cordia africana* Lam., and *Milletia ferruginea* Hochst. Baker are used for shade, fodder, and timber (Kassa et al., 2019).

The general lithology of the area is comprised of underlying Precambrian basement rock overlain by Tertiary volcanic rocks including rhyolites (Schluter et al., 2008; Kassa et al., 2017a; Kassa et al 2017b). Soils are predominantly umbric nitisols on hill slopes and dystic leptosols on crests (Jones et al., 2013). Climate is characterised by unimodal rainfall with the rainy season running from mid-March to mid-November (Kassa et al., 2019). The annual mean rainfall for the nearby town of Mizan Teferi is approximately 1,780 mm (Kassa et al., 2019). The lower sections of the northern slopes of Mount Karkarha averaged ca. 1,313-1,566 mm across three temporary rain gauge stations between 2013 and 2014 (Kassa et al., 2019).

Conservation issues

Mount Karkarha does not fall within a protected area, biodiversity management plan, Key Biodiversity Area, or Important Bird Area

(Protected Planet, 2021).

Satellite imagery from 1984 suggests that forest cover was once more extensive at ca. 18,800 ha, compared to the current ca. 13,757 ha. Forest fragmentation is evident as once connected patches of forest are now disjunct from the current IPA (Google Earth, 2021). This is supported by the findings of Kassa et al. (2017b & 2018) through local knowledge on land use changes. Areas of cropland and agroforestry surrounding the IPA were reportedly natural forest 14-23 years ago (Kassa et al., 2017b). The largest areas of forest loss over the last 30 years have occurred to the south of the IPA where the towns Fachi and Show Gimira have expanded, where areas that were once forest are now agricultural fields or settlement (Google Earth, 2021).

The presence of secondary vegetation, such as *Albizia gummifera*-dominated forest in the northern slopes of the IPA, could suggest historical logging followed by regrowth (Kassa et al., 2019).

Conversion of forest to other land uses is inferred to be a continuing and future threat due to the IPA being encompassed by agricultural fields and bordered by Mizan Teferi to the north and smaller towns and settlements to the east, south, and west.

Chanie & Yirsaw (2018) reported that no natural forest remains in the Mizan-Aman woreda but rather 320 ha of private forest, 19 ha of community forest, and 54 ha of government forest, it is unclear if or how this relates to the forest of Mount Karkarha.

Site assessor(s)

Eden House, Royal Botanic Gardens, Kew
Iain Darbyshire, Royal Botanic Gardens, Kew
Sebsebe Demissew, Addis Ababa University
Sileshi Nemomissa, Addis Ababa University
Ermias Lulekal, Addis Ababa University
Tesfaye Awaw, Ethiopian Biodiversity Institute
Birhanu Belay Telake, Gullele Botanic Garden

IPA criterion A species

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Dorstenia soerensenii</i> Friis	A(i)	✓	✓	✓	—	—	
<i>Scadoxus nutans</i> (Friis & I.Björnstad) Friis & Nordal	A(i)	✓	✓	✓	—	—	

IPA criterion C qualifying habitats

HABITAT	QUALIFYING SUB-CRITERION	≥ 5% OF NATIONAL RESOURCE	≥ 10% OF NATIONAL RESOURCE	1 OF 5 BEST SITES NATIONALLY	AREAL COVERAGE AT SITE
---------	--------------------------	---------------------------	----------------------------	------------------------------	------------------------

General site habitats

GENERAL SITE HABITAT	PERCENT COVERAGE	IMPORTANCE
Forest - Subtropical/Tropical Moist Montane Forest	—	Major
Artificial - Terrestrial - Subtropical/Tropical Heavily Degraded Former Forest	—	Minor

Land use types

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
Harvesting of wild resources	—	Unknown

Threats

THREAT	SEVERITY	TIMING
--------	----------	--------

Management type

MANAGEMENT TYPE	DESCRIPTION	YEAR STARTED	YEAR FINISHED
No management plan in place		—	—

Bibliography

Jones, A., Breuning-Madsen, H., Brossard, M., Dampha, A., Deckers, J., Dewitte, O., Gallali, T., Hallett, S., Jones, R., Kilasara, M., Le Roux, P., Micheli, E., Montanarella, L., Spaargaren, O., Thiombiano, L., Van Ranst, E., Yemefack, M. & Zougmore R. 2013. **Soil Atlas of Africa**.

Protected Planet 2020. **Protected Areas (WDPA)**.

Chanie, M. & Yirsaw, T. 2018. **Economic Contribution of Forest Resources to Sustainable Rural Livelihoods in Bench Maji Zone, South West Ethiopia**.. International Journal of Advanced Research, Vol 6(6), page(s) 1-10

Kassa, H., Dondeyne, S., Poesen, J., Frankl, A., & Nyssem, J. 2017a. **Transition from forest-based to cereal-based agricultural systems: a review of the drivers of land use change and degradation in southwest Ethiopia**.. Land Degradation & Development, Vol 28, page(s) 431-499

Kassa, H., Dondeyne, S., Poesen, J., Frankl, A., & Nyssen, J. 2017b. **Impact of deforestation on soil fertility, soil carbon and nitrogen stocks: the case of the Gacheb catchment in the White Nile Basin, Ethiopia**.. Agriculture, Ecosystems and Environment, Vol 247, page(s) 273-282

Kassa, H., Dondeyne, S., Poesen, J., Frankl, A., & Nyssem, J. 2018. **Agro-ecological implications of forest and agroforestry systems conversion to cereal-based farming systems in the White Nile Basin, Ethiopia**.. Agroecology and Sustainable Food Systems, Vol 42(2), page(s) 149-168

Kassa, H., Frankl, A., Dondeyne, S., Poesen, J., & Nyssen, J. 2019. **Sediment yield at southwest Ethiopia's forest frontier**.. Land Degradation & Development, Vol 30, page(s) 695-705

NTFP 2004. **Baseline description of project area: Summary of participatory appraisal data at Kebele and Got level**. Student Research Series No. 1..

Roux, J.P. 2011. **The fern genus Elaphoglossum section Lepidoglossa (Dryopteridaceae) in Africa, Macaronesia, the mid-Atlantic and southern Indian Ocean Islands**.. Botanical Journal of the Linnean Society, Vol 165, page(s) 20-63

Schluter, T. 2008. **Geological atlas of Africa: with notes on stratigraphy, tectonics, economic geology geohazards and geosites of each country**..

Mesfin Tadesse 2004. **Asteraceae (Compositae)**. In *Flora of Ethiopia and Eritrea*, vol. 4(2), (eds.) Inga Hedberg, Ib Friis, & Sue Edwards..