


Moyen Bafing National Park

National du Moyen Bafing (Test version)

GUITIPA013



Country: **Guinea**

Administrative region: **Dingiraye (Prefecture)**

Central co-ordinates: **11.54056 N, -11.23417 E**

Area: **6426.17km²**

Qualifying IPA criteria

A(i), B(ii), C(iii)

IPA assessment rationale

The proposed Moyen Bafing National Park (MBNP) has 19 threatened species present across different vegetation types. It is the only known location where two Global endemic species are known to survive (*Barleria asterotricha* and *Dissotis linearis*) and has important populations of the globally threatened species *Lipotriche felcis*, *Cyathula pobeguini*, *Macropodiella garrettii*, *Danthoniopsis chevalieri* and *Leocus pobeguini* for which Moyen Bafing contains the only known records (or the only known recent records). Four potential new species have been identified in 2018, all of which are likely to prove critically endangered and potentially globally endemic to Moyen Bafing. In addition, three additional new records for Guinea were discovered. The landscape is under severe pressure from the large local population within the park and the sizeable herds of cattle.

Site description

The proposed Moyen Bafing National Park (MBNP) is in the north-east of the Fouta Djallon, straddling the prefectures of Tougué and Dingiraye. The area extends southwards from the Mali border and encompasses six Classified Forests. It covers a large section of the Bafing river catchment, tributaries of the Falémé river, as well as the Koukoutamba and Téné rivers in the southern part of the park. It is a matrix of vegetation types, including lateritic bowal (bauxite based), wooded grassland, and woodland. It has an altitudinal range of 500m and the highest point (935m) is in the south of the park, at Kourousansan. The rivers have a number of cascades and rapid sections which are home to rare Podostemaceae. It has been designated largely for the large population of Chimpanzees (a population of 3,715 mature individuals) present, as well as other large mammals found predominantly in gallery forest and woodland within the park (listed as priority vegetation types for management in the park). There are 233 villages, 7 hamlets, and 15 camps within the provisional park boundary. This area is made up of three protected zones (ZIP, ZGR, and ZD).

Botanical significance

Three botanical expeditions to the MBNP in 2018 have shown that, despite the fragmented and highly disturbed nature of the vegetation, there are 19 threatened species present, found across a

range of habitat types. In the rapids and cascades, five species of Podostemaceae have been found. In addition, the park has the largest global populations of *Barleria asterotricha* (EN) and *Dissotis linearis* (EN), both endemic to Guinea. A recent collection of *Lipotriche felicis* (EN) was the first collection in Guinea since 1937, and only the third collection globally. There were four potential new species and three new records for Guinea identified in November 2018. The whole park has not been surveyed, so it is possible that more threatened species could be discovered with further surveys.

Camille Reider, Wild Chimpanzee Foundation (WCF)

Habitat and geology

The geology is mostly sandstone, argillite interlayered with limestone, and silica with igneous intrusions. It has a matrix of grassland including lateritic bowal (mostly bauxite based), wooded grassland, and woodland, with small areas of gallery forest. Several large rivers run through the park (Bafing, Koukoutamba, Téné – tributaries of the Falémé in the north and the Bakoun and Kokoun in the centre), with many tributaries and temporary rivers feeding into them. The length of the hydrological network within the park is 4,214km, including 148km of the Bafing River.

Conservation issues

Due to the large human population (c. 67,000 people) within the park, there are significant threats from clearance for fields and cattle grazing. In addition, there are small-scale mining activities and the setting of uncontrolled fires. There are large herds of Ndama cattle across the park and this leads to burning of the grassland for pasture. The increased frequency of burning has almost certainly influenced the species diversity. Some of the bowal areas are also threatened from gravel extraction on a local scale and are also potential sites of mining activities (bauxite). The alluvial areas are also potential sites for gold panning. The potential construction of a hydroelectric dam at the Koukoutamba Falls on the Bafing river (a tributary of the Senegal river) will cause considerable damage to both the populations of threatened Podostemaceae species and threatened habitats. It is predicted that the reservoir for the dam will cover 3.5% of the southern part of the park. The park itself will protect 24% of the hydrological network that will feed the hydroelectric dam.

Site assessor(s)

Assessed by:

Charlotte Couch, Royal Botanic Gardens Kew

Denise Molmou, Herbar National de Guinee/ Simfer

Arnaud Gotanegre, Wild Chimpanzee Foundataion

Martin Cheek, Royal Botanic Gardens Kew

Pacifique Kizila, Wild Chimpanzee Foundataion

Melanie Sottocasa, Wild Chimpanzee Foundataion

IPA criterion A species

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Barleria asterotricha</i> Benoist	A(i)	✓	✓	✓	—	—	Frequent
<i>Mesanthemum tuberosum</i> Lecomte	A(i)	✓	✓	✓	—	—	Scarce
<i>Cyathula pobeguinii</i> Jacq.-Fél.	A(i)	✓	✓	✓	—	—	Scarce
<i>Lipotriche felicis</i> (C.D.Adams) D.J.N.Hind	A(i)	✓	✓	✓	—	—	Occasional
<i>Leocus pobeguinii</i> (Hutch. & Dalziel) J.K.Morton	A(i)	✓	✓	✓	—	—	Scarce
<i>Afzelia africana</i> Sm. ex Pers.	A(i)	—	—	—	—	✓	Frequent
<i>Pterocarpus erinaceus</i> Poir.	A(i)	—	—	—	—	✓	Common
<i>Acridocarpus spectabilis</i> (Nied.) Doorn-Hoekm.	A(i)	✓	✓	✓	—	—	Frequent
<i>Dissotis linearis</i> Jacq.-Fél.	A(i)	✓	✓	✓	—	—	Scarce
<i>Khaya senegalensis</i> A.Juss.	A(i)	—	—	—	—	✓	Common
<i>Embelia djalensis</i> A.Chev. ex Hutch. & Dalziel	A(i)	✓	✓	✓	—	—	Scarce
<i>Inversodicraea koukoutamba</i> Cheek	A(i)	✓	✓	✓	—	—	Abundant
<i>Stonesia taylorii</i> C.Cusset	A(i)	✓	✓	✓	—	—	Frequent
<i>Inversodicraea harrisii</i> (C.Cusset) Cheek	A(i)	✓	✓	✓	—	—	Frequent
<i>Lebbiea grandiflora</i> Cheek	A(i)	✓	✓	✓	—	—	Frequent
<i>Pavetta lasioclada</i> (K.Krause) Mildbr. ex. Bremek.	A(i)	✓	—	✓	—	—	Frequent

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Vitellaria paradoxa</i> C.F.Gaertn.	A(i)	—	—	—	—	✓	Common
<i>Macropodiella garrettii</i> (C. H. Wright) C.Cusset	A(iv)	✓	✓	✓	—	—	Frequent
<i>Calophyllum africanum</i> Cheek & Q.Luke	A(i)	—	✓	✓	—	✓	Frequent
<i>Barleria macclaudii</i> Benoist	A(i)	—	✓	✓	—	—	Scarce

IPA criterion C qualifying habitats

HABITAT	QUALIFYING SUB-CRITERION	≥ 5% OF NATIONAL RESOURCE	≥ 10% OF NATIONAL RESOURCE	1 OF 5 BEST SITES NATIONALLY	AREAL COVERAGE AT SITE
Waterfalls and Rapids	C(iii)	—	✓	✓	877
High Altitude Sandstone Bowal Grasslands	C(iii)	—	✓	✓	

General site habitats

GENERAL SITE HABITAT	PERCENT COVERAGE	IMPORTANCE
Forest - Subtropical/Tropical Dry Forest	—	Minor
Grassland - Subtropical/Tropical Seasonally Wet/Flooded Lowland Grassland	—	Major
Grassland - Subtropical/Tropical High Altitude Grassland	—	Major
Wetlands (inland) - Permanent Rivers, Streams, Creeks [includes waterfalls]	—	Major
Wetlands (inland) - Seasonal/Intermittent/Irregular Rivers, Streams, Creeks	—	Minor
Wetlands (inland) - Seasonal/Intermittent Freshwater Marshes/Pools [under 8 ha]	—	Minor

Land use types

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
Nature conservation	—	Major
Agriculture (arable)	—	Minor
Agriculture (pastoral)	—	Major
Industrial development	—	Unknown
Extractive industry	—	Minor

Threats

THREAT	SEVERITY	TIMING
Agriculture & aquaculture - Annual & perennial non-timber crops - Small-holder farming	Medium	Ongoing - stable
Agriculture & aquaculture - Livestock farming & ranching - Small-holder grazing, ranching or farming	High	Ongoing - increasing
Energy production & mining - Mining & quarrying	Medium	Future - planned activity
Energy production & mining - Renewable energy	High	Future - planned activity
Natural system modifications - Fire & fire suppression - Increase in fire frequency/intensity	High	Ongoing - increasing

Protected areas

PROTECTED AREA NAME	PROTECTED AREA TYPE	RELATIONSHIP WITH IPA	AREAL OVERLAP
Moyen Bafing National Park	National Park	protected/conservation area matches IPA	6426

Management type

MANAGEMENT TYPE	DESCRIPTION	YEAR STARTED	YEAR FINISHED
Site management plan in place	The Guinean government authorised the process of creation for the proposed MBNP in 2017, by a ministerial decree. The TIPA is encompassed by the protected area. A 2018-2020 action plan is available and will lead to a development and management plan by the end of 2020.	2018	2020

Bibliography

Couch, C., Magassouba, S., Rokni, S. & Cheek, M. 2018. **Threatened plants species of Guinea-Conakry: A preliminary checklist**. PeerJ Preprints

IUCN 2019. **IUCN Red List**.

Wild Chimpanzee Foundation 2018. **Plan d'action du projet de création du Parc national du Moyen-Bafing. Version 4..**

Couch, C., Cheek, M. & Molmou, D. 2018-2019. **Trois Rapports de terrain préparés pour Wild Chimpanzee Foundation**.