

# Serra Tumbine

Machinjiri (Test version)

**MOZTIPA036**



Country: **Mozambique**

Administrative region: **Zambézia (Province)**

Central co-ordinates: **-16.08730 N, 35.80350 E**

Area: **13.7km<sup>2</sup>**

## Qualifying IPA criteria

A(i)

## IPA assessment rationale

Serra Tumbine qualifies as an IPA under criterion A. Three sub-criterion A(i) species are recorded from this IPA: *Encephalartos gratus* (VU), *Pavetta chapmanii* (VU) and *Streptocarpus leptopus* (EN). Serra Tumbine is particularly important for the lattermost species as one of only two sites globally and the only Mozambican IPA from which *S. leptopus* is known.

## Site description

Serra Tumbine is a mountain in Milange District, Zambézia Province, Mozambique. The peak reaches 1,548 m and the entire mountain is approximately 8 km in diameter (Woolley 1987). Serra Tumbine is considered to be an outlier of the Mulanje massif, a short distance across the border in Malawi, and is also part of the Mulanje-Namuli-Ribaue centre of endemism (Darbyshire et al. 2019). Milange town lies at the western foot of the mountain and there are a small number of residents who live along the 474 road that runs from the east of Serra Tumbine to Milange. Serra Tumbine has a number of aquifers, and it appears from satellite imagery that there is agricultural land associated with the streams that originate on the mountain, particularly around Milange town.

The area of this IPA is 13.7 km<sup>2</sup>. While the dense montane forest is

of conservation significance, only remnants of this habitat remain due to extensive conversion to agriculture. During the Mozambican Civil War, people fleeing the conflict settled on Serra Tumbine and began cultivating crops on the mountain, particularly the north-eastern slopes (Manuel 2007). It is thought that the loss of forested areas from the mountain may have contributed to a catastrophic landslide in 1998 (World Bank 2019). At lower altitudes the habitat is characterised by miombo woodland, although much of this area has also been converted to agriculture and has been excluded from this IPA.

## Botanical significance

Three threatened species have been recorded from Serra Tumbine. There is one Endangered species, *Streptocarpus leptopus*, which is only known from this IPA and neighboring Mount Mulanje in Malawi and is threatened by clearing of its forest habitat at both sites (Richards 2021). Two Vulnerable species are also known from this site: *Encephalartos gratus* (VU), a cycad mainly threatened in Mozambique by annual burning and the resulting decrease in seedling recruitment (Strugnell 2002; Donaldson 2010; Burrows et al. 2018), and *Pavetta chapmanii* (VU), which is only known from six locations within the Mulanje-Namuli-Ribaue chain of mountains (Timberlake 2020). In addition, a Near Threatened species, *Cola mossambicensis*, has also been collected from Serra Tumbine. Many of the collections for these species are historical so it would be highly desirable to confirm the continued presence of each at this site and establish how large the populations are within this IPA. There has not yet been a full botanical inventory of Serra Tumbine but it is possible that more species of conservation significance are present but have yet to be documented from the site or assessed for the IUCN Red List. For instance, nearby Mount Mulanje is known for its high number of endemic and near endemic species (Strugnell 2002) and, although many of these species are found in high altitude

grasslands and rocky outcrops that are largely absent from Serra Tumbine, we could still expect some species found on Mulanje to be shared with Serra Tumbine.

While the remaining area of mid-altitude forest is not extensive enough for Serra Tumbine to meet C(iii) of the IPA criteria, it has been suggested that the high root density of the forest stabilises the soil and so prevents erosion and landslides (Manuel 2007). Although much of this forest has been cleared for agriculture (World Bank 2019), it is likely that the remaining fragments still provide this important ecosystem service.

---

## Habitat and geology

The remaining mid-altitude montane forest on Serra Tumbine is of great conservation importance. Although only very limited botanical survey work has been conducted here, in a botanical collection at this site by Correia (MF #510), the forest was described as dense, with species including *Albizia*, *Newtonia* (described on the specimen voucher as *Piptadenia* by Correia, but this almost certainly a synonym), *Chrysophyllum* and *Macaranga*. In the nearby Chisongeli forest on Mount Mulanje, *Newtonia buchananii* has been recorded as a dominant species (Dowsett-Lemaire 1988) and it is therefore also likely to be dominant in the forests of Serra Tumbine. There is around 2 – 4 km<sup>2</sup> of forest remaining on Tumbine, mostly located at altitudes above 1,000 m, with the largest patch overlooking Milange town and some smaller patches in steep gullies and on the northerly peak.

On all of the lower slopes of the mountain (On Mount Chipero, forest clearance has been observed to promote the establishment of edge species, such as *Albizia gummifera*, which in turn prevents the re-establishment of forest species (Timberlake et al. 2007). It is possible that the same process has also occurred on Serra Tumbine where forest has been cleared by anthropogenic disturbance or possibly due to landslides.

In terms of geology, Serra Tumbine is a Late Cretaceous to Early Jurassic syenite intrusion within the surrounding Pre-Cambrian metamorphic granulites and gneisses (Woolley 1987; Manuel 2007). The soils derived from the syenites are dark brown with a humic top layer, while the granulites and gneisses form lateritic soils. Both soil types are deep with an overlying layer of colluvial materials varying in size from fine sediments to large boulders (Manuel 2007). Rainfall on the mountain is around 1,200 to 2,000 mm per year, peaking between January and March (Manuel 2007). There is no temperature data for the mountain itself, although nearby Milange town experiences its highest average temperature of 27°C in October and November and an average low of 19°C in June and July (World Weather Online 2021), although it is likely cooler on the upper slopes of the mountain and may possibly experience mists as observed on Mount Chipero to the south (Timberlake et al. 2007).

---

## Conservation issues

This IPA does not fall within a protected area, Important Bird Area (IBA) or Key Biodiversity Area. There has been little scientific

research into the animals of Serra Tumbine, however, the nearby Mount Mulanje is an IBA and it is possible that some of the important bird species from Mulanje occupy Serra Tumbine at least transiently.

As a result of the Mozambican Civil War, people from various places fleeing conflict settled on Serra Tumbine and began occupying the slopes of the mountain and cultivating crops. The lower root density of these crops, compared to the forest that previously occupied these slopes, provides less soil stabilisation and is believed to have contributed to a catastrophic landslide at this site in 1993 (Manuel 2007). These agricultural practices on the slopes of the mountain have continued since the last landslide (Achar 2012).

Other causes of the 1998 landslide are also linked to clearing of vegetation, including high levels of tree felling for charcoal production and fires (Manuel 2007). It is unclear whether the fires have increased in frequency due to anthropogenic burning. Fires are used on Mount Mulanje by hunters to clear the bush (Wisborg & Jumbe 2010) and on Mount Chipero both to hunt and to clear areas for small-scale cultivation (Timberlake et al. 2007); we may therefore expect that at least some of the fires on Serra Tumbine have been related to human activities.

The landslides themselves, of which there have been four between 1940 and 2000, have likely caused massive disturbance to vegetation, as records of the 1993 landslide suggest that high volumes of debris were carried down the mountain (World Bank 2019). This most recent landslide had a huge impact on the local landscape - it is reported that 1,000 hectares of crops were destroyed. The extent of damage to the mid-altitude forest stands is not well-documented, however, it is known that tree trunks came down the mountain in the debris and that large landslides have the potential to clear forests, remove topsoil and make land less productive (Forbes & Broadhead 2013), the latter consequence possibly exacerbating the problem of conversion of forest to agriculture on the mountain.

Support for sustainable food and timber production is required to prevent further landslide catastrophes, which in 1998 caused devastating loss of life and casualties, while also enabling local people to meet their consumption needs. A project run by NGOs Legado and Nitidæ on Mount Namuli has been working with communities to establish a community protected area, secure land rights for local people and promote sustainable economic development for these communities. Part of this work includes a process of delineation a core zone, where conservation is a priority, while using agroecological research to increase production of crops outside this core zone (Nitidæ 2021). A similar approach on Serra Tumbine could help protect and regenerate forests on the mountain, which would both benefit local communities and the threatened species residing in this habitat.

---

## Site assessor(s)

Sophie Richards, Royal Botanic Gardens, Kew

Iain Darbyshire, Royal Botanic Gardens, Kew

## IPA criterion A species

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Pavetta chapmanii</i> <i>Bridson</i>	A(i)	–	✓	✓	–	–	Unknown
<i>Encephalartos gratus</i> <i>Prain</i>	A(i)	–	✓	✓	–	–	Unknown
<i>Streptocarpus leptopus</i> <i>Hilliard &amp; B.L.Burt</i>	A(i)	✓	✓	✓	–	–	Unknown

## IPA criterion C qualifying habitats

HABITAT	QUALIFYING SUB-CRITERION	≥ 5% OF NATIONAL RESOURCE	≥ 10% OF NATIONAL RESOURCE	1 OF 5 BEST SITES NATIONALLY	AREAL COVERAGE AT SITE
Medium Altitude Moist Forest 900-1400 m	C(iii)	–	–	–	1.8

## General site habitats

GENERAL SITE HABITAT	PERCENT COVERAGE	IMPORTANCE
Forest - Subtropical/Tropical Moist Montane Forest	–	Minor
Savanna - Moist Savanna	–	Major
Artificial - Terrestrial - Arable Land	–	Major

## Land use types

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
Agriculture (arable)	–	Major
Harvesting of wild resources	–	Minor

## Threats

THREAT	SEVERITY	TIMING
Agriculture & aquaculture - Annual & perennial non-timber crops - Small-holder farming	High	Ongoing - trend unknown
Natural system modifications - Fire & fire suppression - Increase in fire frequency/intensity	Unknown	Ongoing - trend unknown
Biological resource use - Logging & wood harvesting	Low	Ongoing - trend unknown

## Management type

MANAGEMENT TYPE	DESCRIPTION	YEAR STARTED	YEAR FINISHED
No management plan in place		–	–

## Bibliography

Darbyshire, I., Timberlake, J., Osborne, J., Rokni, S., Matimele, H., Langa, C., Datizua, C., de Sousa, C., Alves, T., Massingue, A., Hadj-Hammou, J., Dhanda, S., Shah, T. & Wursten, B. 2019. **The endemic plants of Mozambique: diversity and conservation status.** PhytoKeys, Vol 136, page(s) 45-96

Burrows, J., Burrows, S., Lötter, M. & Schmidt, E. 2018. **Trees and Shrubs Mozambique.**

Timberlake, J. 2020. **Pavetta chapmanii.** The IUCN Red List of Threatened Species 2020: e.T146652565A146819426.

Timberlake, J., Bayliss, J., Alves, T., Baena, S., Harris, T. & da Sousa, C. 2007. **Biodiversity and Conservation of Mount Chipirone, Mozambique.** Darwin Initiative Award 15/036: Monitoring and Managing Biodiversity Loss in South-east Africa's Montane Ecosystems, page(s) 1-33

Donaldson, J.S. 2010. **Encephalartos gratus.** The IUCN Red List of Threatened Species 2010: e.T41916A10594309.

Achar, J. 2012. **ZAMBÉZIA -Queimadas e Agricultura Rudimentar Podem Provocar Aluimento de Terras Em Gúruè.** MOZAMBIQUE: News reports & clippings

Dowsett-Lemaire, F. 1988. **The Forest Vegetation of Mt Mulanje (Malawi): A Floristic and Chorological Study along an Altitudinal Gradient (650-1950 m).** Bulletin du Jardin botanique national de Belgique / Bulletin van de National Plantentuin van België, Vol 58, page(s) 77-107

Forbes, K. & Broadhead, J. 2013. **Forests and Landslides: The Role of Trees and Forests in the Prevention of Landslides and Rehabilitation of Landslide-Affected Areas in Asia.** Second Edition.

Manuel, I. 2007. **Reduction and Management of Geo-Hazards in Mozambique.** International Journal for Disaster Management & Risk Reduction, Vol 1, page(s) 18-23

Strugnell, A.M. 2002. **Endemics of Mt. Mulanje. The Endemic Spermatophytes of Mt. Mulanje, Malawi.** Systematics and Geography of Plants, Vol 72, page(s) 11-26

World Bank 2019. **Disaster Risk Profile: Mozambique.**

Wisborg, P. & Jumbe, C.B.L. 2010. **Mulanje Mountain Biodiversity Conservation Project: Mid-Term Review for the Norwegian Government.**

Woolley, A.R. 1987. **Alkaline rocks and carbonatites of the world: Africa.**

Nitidæ 2021. **NAMULI SKY ISLAND - Creation of a new protected area around Mount Namuli – Nitidæ.**

Richards, S. 2021. **Streptocarpus leptopus.** The IUCN Red List of Threatened Species.

World Weather Online 2021. **Milange Monthly Climate Averages.**