

the conversion of habitat to agriculture, the presence of *E. zambesiaca* within Sangarassa Forest is not only important for the continued survival of this species but is the only known location within Mozambique and represents the southern edge of its range (Timberlake 2019).

Celosia pandurata (VU), an endemic species, is also recorded from Sangarassa forest. A total of four Mozambican endemics occur within this forest patch and, hence, Sangarassa Forest is of particular importance within this IPA.

Within the IPA as a whole, 12 Mozambican endemics have been recorded. One of these species, *Acacia torrei* (LC) is limited only to a range of ca. 1,700 km² upon the black alluvial clay soils of the Urema and Zangue valleys. This species is locally common in the north of the IPA and is currently assessed as Least Concern; the continued protection *A. torrei* receives within Gorongosa National Park is central to preventing it becoming globally threatened with extinction (Coates Palgrave et al. 2014). The Urema Valley also hosts the largest known population of another endemic species, *Gyrodroma hispida* (LC). *G. hispida* has been described as common within this IPA and so this is a key locality for preventing this species becoming threatened with extinction (Richards 2021).

The over 700 km² of seasonally inundated grasslands within this IPA represent a habitat type of conservation interest for Mozambique (Stalmans & Beilfuss 2008). As well as hosting significant populations of the two endemic species mentioned above, this habitat type has a limited range across Mozambique. The seasonally inundated grasslands of this site represent one of the largest and highest quality examples of this habitat nationally and therefore trigger sub-criterion C(iii) of the IPA criteria for this site.

Habitat and geology

The plant communities within this IPA are highly variable, which likely reflects the underlying soil structure and moisture availability within the landscape (Stalmans & Beilfuss 2008). Much of the IPA has sandy soils, with black clay colluvial fan to the north (Steinbruch 2010). *Acacia torrei* in particular is reliant on these areas of black clay and is restricted to this substrate within Sofala Province (Coates Palgrave et al. 2014).

Lake Urema, just south of the centre of the valley, is supplied with drainage from both Mount Gorongosa and the surrounding plateau to the west and the Cheringoma Plateau to the east, with overflow joining the Pungue River at the southern boundary of this IPA (Stalmans & Beilfuss 2008). During the wet season, December to March, the water levels of Lake Urema and associated rivers increase to cover up to 40% of Gorongosa's area (Stahl 2020), with much of the central stretch of this IPA, along with the southern boundary by the Pungue River, being inundated with water (Parque Nacional da Gorongosa 2019).

The vegetation types of Gorongosa National Park are categorised in Stalmans & Beilfuss (2008) and a summary of the relevant vegetation types is provided below.

The floodplain region is a largely open landscape dominated by

seasonally inundated grasslands of various types including: *Echinochloa* – *Chrysopogon*, *Setaria* and *Cynodon dactylon* – *Digitaria didactyla* assemblages; the latter community is concentrated around Lake Urema and has almost no woody plants. On the lower slopes and drainage lines south of Lake Urema are areas of palm savanna consisting of open to closed *Hyphaene* stands with a grassy understory. Stands of *Acacia xanthophloea*, mixed *Acacia-Combretum* and *Faidherbia albida* also form open to closed areas of woodland within the floodplain and alluvial fan. West of Chitengo Camp is Sangarassa Forest, a 1.6 km² area of vegetation that is described on specimen vouchers as dense sand forest (e.g., Wursten #911). The forest is dominated by species such as *Newtonia hildebrandtii* and *Xylocarpus torreana* (Tinley #2331). The understory includes the Mozambican endemic *Millettia mossambicensis*, while some species are associated with the termite mounds that border seasonal pans, such as *Cola mossambicensis* (NT).

To the south of the IPA, following the Pungue River, is closed woodland/dry forest dominated by *Piliostigma thonningii* and, in seasonally flooded areas, *Borassus aethiopicum* (Stalmans & Beilfuss 2008; Hyde et al. 2020a, 2020b).

Conservation issues

The entirety of the Urema Valley and Sangarassa Forest falls within Gorongosa National Park and Buffer Zone with only the most northerly 220 km² of the IPA falling within the buffer zone. This IPA is also encompassed within Gorongosa Mountain and National Park Important Bird Area and Gorongosa-Marromeu Key Biodiversity Area. The Urema Valley wetlands are of particular importance for bird species; Grey Crowned Crane (*Balearica regulorum*- EN) has been recorded here, while the area is possibly an important over-wintering ground for Great Snipe (*Gallinago media*- NT) (BirdLife International 2020). In addition, a 2014 count found that the population sizes of two avian taxa meet Ramsar site criteria; Yellow-billed Stork (*Mycteria ibis*- LC) exceeded the threshold of 1% of the population in sub-Saharan Africa (870 individuals) and the population of African Darter (*Anhinga rufa*- LC) exceeded 1% of individuals of this species in southern and eastern Africa (1,000 individuals) (Stalmans et al. 2014), although the area is not currently listed as a Ramsar site.

Mimosa pigra, a species that features in IUCN's "100 of the World's Worst Invasive Alien Species" (van der Weijden et al. 2004), is a major threat to the wetlands within this IPA and has established on the floodplain (Stalmans & Beilfuss 2008). This species forms dense thickets, excluding other species and converting floodplains into scrubland (Beilfuss 2007). However, it is thought that the re-introduction of ruminant grazers is helping to contain shrub encroachment (Guyton et al. 2020).

The management strategy for the site includes a cold burning of the valley early in the dry season (Stahl 2020). Research into fire and herbivory dynamics has been undertaken (see Stahl 2020) towards improving the use of fire for the continued restoration of the park following the Mozambican Civil War.

During the civil war, large herbivore populations declined by over 90% within Gorongosa National Park (Stalmans et al. 2019). Today, as populations continue to recover, a number of species within the national park are centred around the Urema Valley region. Hippo were released around Lake Urema in 2008 and African wild dog released in 2018, while the recovering Sungwe lion pride is centred around the streams south-west of Lake Urema. The vegetation within this IPA, as an important component of the ecosystem, therefore, makes an important contribution to the conservation of mammals within the national park, as well as the tourism that these mammal species attract.

However, as the number of large herbivores within the park increases, a change in dominant species has been recorded. Elephant, hippo and African buffalo previously dominated the pre-war large herbivore biomass, while in 2018 over 74% of large herbivore biomass recorded was waterbuck (Stalmans et al. 2019). It would be informative to monitor how these changing herbivory dynamics may be impacting plant communities, particularly habitats vital for rare or threatened plants species.

Much of this IPA has been unaffected by conversion of habitat to agriculture, a major threat to plant species across Mozambique (Darbyshire et al. 2019), likely because the vast majority of the IPA area falls within Gorongosa National Park. A total of ca. 200,000 people live within the buffer zone and Gorongosa National Park partners with these communities to build sustainable livelihood opportunities (Parque Nacional da Gorongosa 2019). To this end, GNP are working towards having a large Community Conservation Area proclaimed at the north-east boundary, which would include part of this IPA (M. Stalmans, pers. comm. 2021). Monitoring of populations of *Acacia torrei* in this area could be considered within conservation actions, to safeguard against threats to a key area of habitat for this range restricted endemic.

Site assessor(s)

Sophie Richards, Royal Botanic Gardens, Kew

Iain Darbyshire, Royal Botanic Gardens, Kew

IPA criterion A species

SPECIES	QUALIFYING SUB-CRITERION	≥ 1% OF GLOBAL POPULATION	≥ 5% OF NATIONAL POPULATION	1 OF 5 BEST SITES NATIONALLY	ENTIRE GLOBAL POPULATION	SOCIO-ECONOMICALLY IMPORTANT	ABUNDANCE AT SITE
<i>Erythrococca zambesiaca</i> Prain	A(i)	✓	✓	✓	—	—	Unknown
<i>Celosia pandurata</i> Baker	A(i)	✓	✓	—	—	—	Unknown
<i>Vepris myrei</i> (Exell & Mendonça) Mziray	A(i)	✓	✓	✓	—	—	Frequent

IPA criterion C qualifying habitats

HABITAT	QUALIFYING SUB-CRITERION	≥ 5% OF NATIONAL RESOURCE	≥ 10% OF NATIONAL RESOURCE	1 OF 5 BEST SITES NATIONALLY	AREAL COVERAGE AT SITE
Seasonally Inundated Grassland	C(iii)	—	—	—	780

General site habitats

GENERAL SITE HABITAT	PERCENT COVERAGE	IMPORTANCE
Grassland - Subtropical/Tropical Seasonally Wet/Flooded Lowland Grassland	—	Major
Savanna - Moist Savanna	—	Major
Forest - Subtropical/Tropical Dry Forest	—	Minor

Land use types

LAND USE TYPE	PERCENT COVERAGE	IMPORTANCE
Nature conservation	100	Major
Tourism / Recreation	—	Major
Agriculture (arable)	—	Minor

Threats

THREAT	SEVERITY	TIMING
Climate change & severe weather - Storms & flooding	High	Past, likely to return
Natural system modifications - Fire & fire suppression - Trend Unknown/Unrecorded	Unknown	Ongoing - stable
Agriculture & aquaculture - Annual & perennial non-timber crops - Small-holder farming	Low	Ongoing - trend unknown

Protected areas

PROTECTED AREA NAME	PROTECTED AREA TYPE	RELATIONSHIP WITH IPA	AREAL OVERLAP
Gorongosa National Park and Buffer Zone	National Park	protected/conservation area encompasses IPA	1280

Conservation designation

DESIGNATION NAME	PROTECTED AREA	RELATIONSHIP WITH IPA	AREAL OVERLAP
Gorongosa Mountain and National Park	Important Bird Area	protected/conservation area overlaps with IPA	1030
Gorongosa-Marromeu	Key Biodiversity Area	protected/conservation area encompasses IPA	1280

Management type

MANAGEMENT TYPE	DESCRIPTION	YEAR STARTED	YEAR FINISHED
Protected Area management plan in place	The focus of Gorongosa has been set out in the 2020-2050 Strategic Plan and involves improving the capacity of the national park to "preserve, protect and manage the diverse ecosystems within the Park" while also working with communities within the buffer zone, making a particular effort to reach women in these communities, to improve sustainable economic opportunities (Parque Nacional de Gorongosa 2019).	2020	2050

Bibliography

BirdLife International 2021. **Important Bird Areas factsheet: Gorongosa Mountain and National Park.**

Darbyshire, I., Timberlake, J., Osborne, J., Rokni, S., Matimele, H., Langa, C., Datizua, C., de Sousa, C., Alves, T., Massingue, A., Hadj-Hammou, J., Dhanda, S., Shah, T. & Wursten, B. 2019. **The endemic plants of Mozambique: diversity and conservation status.** *PhytoKeys*, Vol 136, page(s) 45-96

Coates Palgrave, F.M., Burrows, J.E., Timberlake, J., Alves, M.T., Contu, S., Hyde, M.A., Luke, W.R.Q., Massingue, A.O., Matimele, H.A., Raimondo, D., Osborne, J. & Hadj-Hammou, J. 2014. **Acacia torrei.** *The IUCN Red List of Threatened Species 2014: e.T19891788A63707954.*

Parque Nacional da Gorongosa 2019. **Our Gorongosa - A Park for the People. Annual Report 2019..**

Quammen, D. 2019. **How one of Africa's great parks is rebounding from war.** *National Geographic*

Stalmans, M., Beilfuss, R. 2008. **Landscapes of Gorongosa National Park.**

Beilfuss, R. 2007. **Adaptive Management of the Invasive Shrub *Mimosa pigra* at Gorongosa National Park.**

Guyton, J.A., Pansu, J., Hutchinson, M.C., Kartzinel, T.R., Potter, A.B., Coverdale, T.C., Daskin, J.H., da Conceição, A.G., Peel, M.J.S., Stalmans, M.E., & Pringle, R.M. 2020. **Trophic rewilding revives biotic resistance to shrub invasion.** *Nature Ecology and Evolution*, Vol 4, page(s) 5

Hyde, M. A., Wursten, B. T., Ballings, P., & Coates Palgrave, M. 2020a. **Flora of Mozambique: Google Maps: *Borassus aethiopum*.**

Hyde, M. A., Wursten, B. T., Ballings, P., & Coates Palgrave, M. 2020b. **Flora of Mozambique: Google maps: *Piliostigma thonningii*.**

Stahl, M. 2020. **Pyric herbivory: Understanding fire-herbivore interactions in Gorongosa National Park..**

Stalmans, M.E., Massad, T.J., Peel, M.J.S., Tarnita, C.E., & Pringle, R.M. 2019. **War-induced collapse and asymmetric recovery of large-mammal populations in Gorongosa National Park, Mozambique.** PLoS ONE, Vol 14

Stalmans M., Davies G.B.P., Trollip J. & Poole G. 2015. **A Major Waterbird Breeding Colony at Lake Urema, Gorongosa National Park, Moçambique.** Durban Natural Science Museum Novitates, Vol 37, page(s) 54-57

Steinbruch, F. 2010. **Geology and geomorphology of the Urema Graben with emphasis on the evolution of Lake Urema.** Journal of African Earth Sciences, Vol 58, page(s) 272-284

van der Weijden, W., Leewis, R., & Bol, P. 2004. **100 of the World's Worst Invasive Alien Species: A selection from the Global Invasive Species Database.** Biological Globalisation, page(s) 206–208

Vieira, B.M. 2011. **A preliminary study on the distribution of *Mimosa pigra* in Gorongosa National Park: reasons and causes.**

Warren, M. 2019. **Why Cyclone Idai is one of the Southern Hemisphere's most devastating storms.** Nature News

Timberlake, J. 2019. ***Erythrococca zambesiaca*.** The IUCN Red List of Threatened Species 2019: e.T146427908A146819180.